

# RADIESSE®

**Boost your  
skin's radiance<sup>1,2</sup>  
and firmness<sup>2-4</sup>  
on your GLP-1  
weight loss  
journey**

**Ask your injector about  
RADIESSE® today**

Firm and tighten your skin while  
reversing visible signs of aging.<sup>1,3,5-10</sup>

Actual patient.  
Results may vary.



## **What are RADIESSE® and RADIESSE® (+)?**

RADIESSE® and RADIESSE® (+) are dermal fillers that are used for smoothing moderate to severe facial wrinkles and folds, such as nasolabial folds (the creases that extend from the corner of your nose to the corner of your mouth). RADIESSE® is also used for correcting volume loss in the back of the hands. RADIESSE® (+) is also used for improving moderate to severe loss of jawline contour in adults over 21 years old.

Please see **Important Safety Information** for RADIESSE and RADIESSE (+) on the next page, and visit [www.RADIESSE.com](http://www.RADIESSE.com) for RADIESSE/RADIESSE (+) **Instructions For Use**.

# RADIESSE®

## Regenerate skin health

### 96%

more elastin  
than baseline<sup>11\*</sup>

### 94%

of patients showed improved  
smiles lines than baseline<sup>5†</sup>

### 98%

patient  
satisfaction<sup>12‡</sup>

\*Based on 6-month human tissue study vs placebo.

†In a randomized split-face study (n=117), 94% of calcium hydroxylapatite-treated folds were rated as improved at 6 months (Global Aesthetic Improvement Scale).

‡In a multicenter, randomized study (n=120), 98% of calcium hydroxylapatite-treated patients self-reported overall satisfaction at 4 months after the second injection.

### RADIESSE® and RADIESSE® (+) IMPORTANT CONSUMER SAFETY INFORMATION

#### Who should not use RADIESSE® or RADIESSE® (+)?

You should not use RADIESSE® or RADIESSE® (+) if you have an allergy to any component of the product, if you have a history of severe allergies, if you have a bleeding disorder, or if you are pregnant or breastfeeding. You should not use RADIESSE® (+) if you have an allergy to lidocaine or medicines like it.

#### What is the most important information I should know about RADIESSE® and RADIESSE® (+)?

One of the risks with using these products is unintentional injection into a blood vessel. The chances of this happening are very small, but if it does happen, the complications can be serious, and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring of the skin. If you have changes in your vision, signs of a stroke (including sudden difficulty speaking, numbness or weakness in your face, arms, or legs, difficulty walking, face drooping, severe headache, dizziness, or confusion), white appearance of the skin, or unusual pain during or shortly after treatment, you should notify your health care practitioner immediately.

As with all procedures that involve an injection through the skin, there is a risk of infection.

Do not use RADIESSE® or RADIESSE® (+) if you have a skin infection until it has healed.

It is not known if RADIESSE® or RADIESSE® (+) is safe or effective in the lips, or in the area around the eyes.

It is not known if RADIESSE® or RADIESSE® (+) is safe or effective beyond 3 years in the face and 1 year in the hand.

Injection of RADIESSE® (+) into the jawline may temporarily alter jaw function.

The microspheres in RADIESSE® and RADIESSE® (+) can be seen in X-rays and CT Scans. It is very important that you tell your health care provider that you have had RADIESSE® or RADIESSE® (+) dermal filler.

If you have a history of herpes, you may experience a herpes breakout after receiving RADIESSE® or RADIESSE® (+).

Injection in the back of the hand may result in temporary difficulty performing activities. RADIESSE® may cause nodules, bumps or lumps in the back of the hand and can last up to 1 year.

You should minimize strenuous activity and avoid extensive sun or heat exposure for about 24 hours after treatment and until any swelling or redness has resolved.

#### What should I tell my doctor before using RADIESSE® or RADIESSE® (+)?

Tell your health care provider if you are taking blood thinners or medicines that can interfere with the clotting of blood, such as

aspirin or warfarin. These medicines might make it more likely that you will experience bruising or bleeding at the injection site. Tell your health care provider if you have any diseases, injuries or disabilities of the hand, if you have a history forming large, raised scars or if you have had any other skin treatments such as skin peels.

#### What are the most common adverse events with RADIESSE® or RADIESSE® (+)?

The most common adverse events seen in clinical studies of RADIESSE® used in the hands include bruising, redness, swelling, pain, itching, nodules or bumps/lumps, difficulty performing activities, loss of sensation and other local side effects. The most common adverse events seen in clinical studies of RADIESSE® or RADIESSE® (+) used in the face include bruising, redness, swelling, pain, itching, difficulty chewing, and other local side effects.

Delayed-onset inflammation near the site of injection is one of the known adverse events associated with dermal fillers. Cases of delayed-onset inflammation have been reported to occur at the treatment site following viral or bacterial illnesses or infections, vaccinations, or dental procedures. Typically, the reported inflammation was responsive to treatment or resolved on its own.

These are not all of the possible side effects with RADIESSE® or RADIESSE® (+). Merz collects information about adverse events seen with RADIESSE® and RADIESSE® (+) outside of clinical studies. These events are included in the RADIESSE® and RADIESSE® (+) Patient Information Guide based on an assessment of seriousness and potential causal relationship to RADIESSE® or RADIESSE® (+). Please see the Patient Information Guide available at [www.radiesse.com](http://www.radiesse.com) for list of these events. Tell your health care provider about any side effects that bother you or do not go away.

Important: For full safety information, please visit [www.Radiesse.com](http://www.Radiesse.com) or call MyMerz Solutions at 1-844-469-6379

RADIESSE® and RADIESSE® (+) are available by prescription only.

**References:** 1. González N, et al. *Dermatol Surg.* 2019;45(4):547-551. 2. Aguilera SB, et al. *Aesthet Surg J.* 2023;43(10):1063-1090. 3. Yutskovskaya YA, et al. *J Drugs Dermatol.* 2017;16(1):68-74. 4. Yutskovskaya YA, et al. *J Drugs Dermatol.* 2020;19(4):405-411. 5. Bass LS, et al. *Aesthet Surg J.* 2010;30(2):235-238. 6. Cogorno Wasylkowski V. *Clin Cosmet Investig Dermatol.* 2015;8:267-273. 7. Draoui O. Chapter 1: The Story of Calcium Hydroxylapatite. In: van Loghem J, ed. *Calcium Hydroxylapatite Soft Tissue Fillers, Expert Treatment Techniques*. 1st ed. Taylor & Francis Group; 2020. 8. Kim J, et al. *Clin Cosmet Investig Dermatol.* 2019;12:771-784. 9. Silvers SL, et al. *Plast Reconstr Surg.* 2006;118(3 Suppl):34S-45S. 10. Moradi A, et al. *J Drugs Dermatol.* 2021;20(11):1231-1238. 11. Data on file. Merz North America, Inc. 2025. 12. Moers-Carpi M, et al. *Dermatol Surg.* 2007;33(suppl 2):S144-S151.

## RADIESSE® | MERZ AESTHETICS®

© 2025 Merz North America, Inc. All rights reserved. MERZ AESTHETICS is a registered trademark of Merz Pharma GmbH & Co. KGaA in the U.S. RADIESSE is a registered trademark of Merz North America, Inc. in the U.S. US-RAD-2500077